

Gauge-invariant Non-spherical Metric Perturbations of Schwarzschild Black-Hole Spacetimes

Alessandro Nagar^{†*} and Luciano Rezzolla^{‡§}

[†]Department of Astronomy and Astrophysics, University of Valencia, Burjassot, Spain

^{*}Department of Physics at the Politecnico, Turin, Italy

[‡]SISSA, International School for Advanced Studies and INFN, Trieste, Italy

[§]Department of Physics, Louisiana State University, Baton Rouge, USA

Abstract. The theory of gauge-invariant non-spherical metric perturbations of Schwarzschild black hole spacetimes is now well established. Yet, as different notations and conventions have been used throughout the years, the literature on the subject is often confusing and sometimes confused. The purpose of this paper is to review and collect the relevant expressions related to the Regge-Wheeler and Zerilli equations for the odd and even-parity perturbations of a Schwarzschild spacetime. Special attention is paid to the form they assume in the presence of matter-sources and, for the two most popular conventions in the literature, to the asymptotic expressions and gravitational-wave amplitudes. Besides pointing out some inconsistencies in the literature, the expressions collected here could serve as a quick reference for the calculation of the perturbations of Schwarzschild black hole spacetimes driven by generic sources and for those approaches in which gravitational waves are extracted from numerically generated spacetimes.

PACS numbers: 04.30.Db, 04.40.Dg, 04.70.Bw,

1. Introduction

Black-hole perturbation theory (see for example the monograph by Chandrasekhar [1], the book by Frolov and Novikov [2] or the recent review by Kokkotas and Schmidt [3]) has been fundamental not only for understanding the stability and oscillations properties of black hole spacetimes [4], but also as an essential tool for clarifying the dynamics that accompanies the process of black hole formation as a result of gravitational collapse [5, 6]. As one example among the many possible, the use of perturbation theory has led to the discovery that Schwarzschild black holes are characterised by decaying modes of oscillation that depend on the black hole mass only, i.e. the black hole quasi-normal modes [7, 8, 9, 10]. Similarly, black-hole perturbation theory and the identification of a power-law decay in the late-time dynamics of generic black-hole perturbations has led to important theorems, such as the ‘no hair’ theorem, underlining the basic black-hole property of removing all perturbations so that ‘all that can be radiated away is radiated away’ [5, 6, 11].

The foundations of non-spherical metric perturbations of Schwarzschild black holes date back to the work in 1957 of Regge and Wheeler [4], where the linear stability of the Schwarzschild singularity was first addressed. A number of investigations, both gauge-invariant and not, then followed in the 1970s, when many different approaches were proposed and some of the most important results about the physics of perturbed spherical and rotating black holes established [5, 6, 7, 8, 10, 12, 13, 14, 15, 16, 17, 18]. Building on these studies, which defined most of the mathematical apparatus behind generic perturbations of black

holes, a number of applications have been performed to study, for instance, the evolutions of perturbations on a collapsing background spacetime [19, 20, 21, 22, 23, 24, 25, 26]. Furthermore, the gauge-invariant and coordinate independent formalism for perturbations of spherically symmetric spacetimes developed in the 70's by Gerlach and Sengupta in [19, 20, 21], has been recently extended to higher-dimensional spacetimes with a maximally symmetric subspace in [27, 28, 29, 30], for the study of perturbations in brane-world models.

Also nowadays, when numerical relativity calculations allow to evolve the Einstein equations in the absence of symmetries and in fully nonlinear regimes, black hole perturbative techniques represent important tools. Schwarzschild perturbation theory, for instance, has been useful in studying the late-time behaviour of the coalescence of compact binaries in a numerical simulation after the apparent horizon has formed [31, 32, 33]. In addition, methods have been developed that match a fully numerical and three-dimensional Cauchy solution of Einstein's equations on spacelike hypersurfaces with a perturbative solution in a region where the components of three-metric (or of the extrinsic curvature) can be treated as linear perturbations of a Schwarzschild black hole (this is usually referred to as the 'Cauchy-Perturbative matching') [34, 35, 36, 37, 38]. This method, in turn, allows to "extract" the gravitational waves generated by the simulation, evolve them out to the wave-zone where they assume their asymptotic form, and ultimately provide outer boundary conditions for the numerical evolution.

Overall, therefore, black hole perturbation theory represents a very powerful tool with multiple applications and developments. The purpose of this review is to review the theory of metric perturbations of Schwarzschild black holes, especially in its gauge-invariant formulations. While a systematic derivation of the most relevant expressions is provided, special care is paid to 'filter' those technical details that may obscure the important results and provide the reader with a set of expressions that can be readily used for the calculation of the odd and even-parity perturbations of a Schwarzschild spacetime in the presence of generic matter-sources. Also, an effort is made to 'steer' the reader through the numerous conventions and notations that have accompanied the development of the formalism over the years, pointing out the misprints and errors that can be found in the literature. Finally, although most of what presented here is not new, some expressions that have not been discussed before in the literature are also introduced.

It should be noted that our discussion will be restricted exclusively to black-hole spacetimes, although a Schwarzschild metric can also be used to describe the vacuum exterior of a spherical relativistic star. The reason behind this choice is that boundary conditions different from the ones used for black holes need to be imposed at the stellar surface. Furthermore, in the case of a relativistic star the Einstein equations need to be suitably coupled with the equations of relativistic hydrodynamics. A detailed discussion of how to do this in practice is beyond the purpose of this review, which is instead focused on a vacuum background spacetime. However, a recent review on the theory of relativistic stellar perturbations can be found in [3].

The organisation of the paper is as follows. In section 2 we review the basic properties and advantages of a gauge-invariant description of the perturbations of a curved spacetime, while in section 3 we review the multipolar expansion of the perturbations of a Schwarzschild spacetime. Sections 4 and 5 are instead devoted to the derivation of the general equations governing the evolution of odd and even-parity perturbations, respectively, whose asymptotic expressions for the gravitational-wave amplitudes and losses are presented in section 6. Finally, section 7 contains our conclusions and a number of Appendices follow to provide some useful relations and expressions for the angular parts of the perturbations.

We use a spacelike signature $(-, +, +, +)$ with Greek indices running from 0 to 3 and

indicate partial and covariant derivatives in the μ -th direction with the symbols ∂_μ and ∇_μ , respectively. Also, symmetrized and antisymmetrized indices in a tensor will be indicated with round and square brackets, respectively.

2. Gauge-invariant metric perturbations

To highlight the importance of gauge-invariant quantities in describing the metric perturbations it is useful to recall that even if the coordinate system of the background spacetime has been fixed, the coordinate freedom of general relativity introduces a problem when linear perturbations are added. In particular, it is not possible to distinguish an infinitesimal ‘physical’ perturbation from one produced as a result of an infinitesimal coordinate transformation (or gauge-transformation). This difficulty, however, can be removed either by explicitly fixing a gauge (as done in the pioneering works on the subject [4, 5, 6, 7, 8, 12, 13]), or by introducing linearly gauge-invariant perturbations (as initially suggested by Moncrief [14] and subsequently adopted in several applications [15, 16, 23, 24, 25, 26]).

More specifically, given a tensor field $\mathbf{X}$ in a background metric $\overset{\circ}{g}$ and $\delta\mathbf{X}$ its infinitesimal perturbation, an infinitesimal coordinate transformation $x^\mu \rightarrow x^{\mu'} \equiv x^\mu + \xi^\mu$ with $\xi^\mu \ll 1$ will yield a new tensor field

$$\delta\mathbf{X} \rightarrow \delta\mathbf{X}' = \delta\mathbf{X} + \mathcal{L}_\xi \mathbf{X} , \quad (1)$$

where $\mathcal{L}_\xi$ is the Lie derivative along ξ in the metric $\overset{\circ}{g}$. We will then consider $\delta\mathbf{X}$ to be gauge-invariant if and only if $\mathcal{L}_\xi \mathbf{X} = 0$.

Stated differently, the possibility of building gauge-invariant metric perturbations relies on the existence of symmetries of the background metric. In the case of a general spherically symmetric background spacetime (i.e. one allowing for a time dependence) and which has been decomposed in multipoles (see Sect. 3), the construction of gauge-invariant quantities is possible for multipoles of order $\ell \geq 2$ only [19, 20, 39, 40]. In practice, the advantage in the use of gauge-invariant quantities is that they are naturally related to scalar observables and, for what is relevant here, to the energy and momentum of gravitational waves. At the same time, this choice guarantees that possible gauge-dependent contributions are excluded by construction.

Of course, this procedure is possible if and only if the background metric has the proper symmetries under infinitesimal coordinates transformation and a gauge-invariant formulation of the Einstein equations for the perturbations of a general spacetime is not possible. Nevertheless, since any asymptotically flat spacetime can in general be matched to a Schwarzschild one at sufficiently large distances, a gauge-invariant formulation can be an effective tool to extract physically information about the gravitational waves generated in a numerically evolved, asymptotically flat spacetime [34, 35, 36, 37, 38].

3. Multipolar expansion of metric perturbations

Given a spherically symmetric Schwarzschild solution with metric $\overset{\circ}{g}$ and line element

$$ds^2 \equiv \overset{\circ}{g}_{\mu\nu} dx^\mu dx^\nu = -e^{2a} dt^2 + e^{2b} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) , \quad (2)$$

where $e^{2a} = e^{-2b} = (1 - 2M/r)$, we generically consider small non-spherical perturbations $h_{\mu\nu}$ such that the new perturbed metric is

$$g_{\mu\nu} \equiv \overset{\circ}{g}_{\mu\nu} + h_{\mu\nu} , \quad (3)$$

where $|h_{\mu\nu}|/|\overset{\circ}{g}_{\mu\nu}| \ll 1$. It should be noted that although we have here chosen to employ Schwarzschild coordinates to facilitate the comparison with much of the previous literature, this is not the only possible choice. Indeed, it is possible to formulate the perturbations equations independently of the choice of coordinates as discussed in [49] and in [51].

Because the background manifold $\mathcal{M}$ is spatially spherically symmetric, it can be written as the product $\mathcal{M} = \mathcal{M}^2 \times S^2$, where $\mathcal{M}^2$ is a Lorentzian 2-dimensional manifold of coordinates (t, r) and S^2 is the 2-sphere of unit radius and coordinates (θ, ϕ) . As a result of this decomposition, the perturbations can be split *ab initio* in a part confined to $\mathcal{M}^2$ and in a part confined on the 2-sphere S^2 of metric γ . Exploiting this, we can expand the metric perturbations h in multipoles referred to as odd or even-parity according to their transformation properties under parity. In particular, are *odd* (or *axial*) multipoles those that transform as $(-1)^{\ell+1}$, under a parity transformation $(\theta, \phi) \rightarrow (\pi - \theta, \pi + \phi)$, while are *even* (or *polar*) those multipoles that transform as $(-1)^\ell$. As a result, the metric perturbations can be written as

$$h_{\mu\nu} = \sum_{\ell, m} \left[(h_{\mu\nu}^{\ell m})^{(o)} + (h_{\mu\nu}^{\ell m})^{(e)} \right], \quad (4)$$

where

$$\sum_{\ell, m} \equiv \sum_{\ell=2}^{\infty} \sum_{m=-\ell}^{\ell}.$$

Introducing now a notation inspired by that of Gerlach and Sengupta [19, 20, 21] and recently revived by Gundlach and Martín-García [39, 40, 41], we use upper-case indices $A, B \dots = 0, 1$ to label the coordinates of $\mathcal{M}^2$ and lower-case indices $c, d \dots = 2, 3$ to label the coordinates of S^2 .

Using this notation, the scalar spherical harmonics $Y^{\ell m}$ are then simply defined as

$$\gamma^{cd} \nabla_d \nabla_c Y^{\ell m} = -\Lambda Y^{\ell m}, \quad (5)$$

where ∇_c indicates the covariant derivative with respect to the metric $\gamma \equiv \text{diag}(1, \sin^2 \theta)$ of S^2 and where $\Lambda \equiv \ell(\ell + 1)$. It is now convenient to express the odd and even-parity metric functions in (3) in terms of tensor spherical harmonics. To do this we introduce the axial vector $S_c^{\ell m}$ defined as

$$S_c^{\ell m} \equiv \epsilon_{cd} \gamma^{de} \nabla_e Y^{\ell m}, \quad (6)$$

where ϵ_{cd} is the volume form on S^2 as defined by the condition $\epsilon_{cd} \epsilon^{ce} = \gamma_d^e$ and such that $\nabla_c \epsilon_{ab} = 0$. In this way, each odd-parity metric function in (3) can be written as

$$(h_{\mu\nu}^{\ell m})^{(o)} = \begin{pmatrix} 0 & h_A^{(o)} S_c^{\ell m} \\ h_A^{(o)} S_c^{\ell m} & h \nabla_{(d} S_{c)}^{\ell m} \end{pmatrix}, \quad (7)$$

where $h_A^{(o)}$, h are functions of (t, r) only and where we have omitted the indices ℓ, m for clarity.

Proceeding in a similar manner, each even-parity metric function can be decomposed in tensor spherical harmonics as

$$(h_{\mu\nu}^{\ell m})^{(e)} = \left(\begin{array}{cc|c} e^{2a} H_0 Y^{\ell m} & H_1 Y^{\ell m} & h_A^{(e)} \nabla_c Y^{\ell m} \\ H_1 Y^{\ell m} & e^{2b} H_2 Y^{\ell m} & \\ \hline h_A^{(e)} \nabla_c Y^{\ell m} & & r^2 (K Y^{\ell m} \gamma_{cd} + G \nabla_d \nabla_c Y^{\ell m}) \end{array} \right), \quad (8)$$

where $H_0, H_1, H_2, h_0^{(e)}, h_1^{(e)} K, G$ (with the indices ℓ, m omitted for clarity) are the coefficients of the even-parity expansion, are also functions of (t, r) only.

Note that we have here used the Regge-Wheeler set of tensor harmonics to decompose the even-parity part of the metric in multipoles [4]. Despite this being a popular choice in the literature, it is not the most convenient one since the tensor harmonics in this set are not linearly independent. An orthonormal set is instead given by the Zerilli-Mathews tensor harmonics [42, 43] and the transformation from one basis to the other is given by defining the tensor $Z_{cd}^{\ell m}$ confined on the 2-sphere S^2

$$Z_{cd}^{\ell m} \equiv \nabla_c \nabla_d Y^{\ell m} + \frac{\Lambda}{2} \gamma_{cd} Y^{\ell m}, \quad (9)$$

and then replacing in equation (8) the second covariant derivative of the spherical harmonics $\nabla_c \nabla_d Y^{\ell m}$ with $Z_{cd}^{\ell m}$. This transformation has to be taken into account, for instance, when developing gauge-invariant procedures for extracting the gravitational-wave content from numerically generated spacetimes which are “almost” Schwarzschild spacetime [44, 45, 46, 47, 48].

Besides vacuum tensor perturbations, the background Schwarzschild spacetime can be modified if non-vacuum tensor perturbations are present and have a mass-energy much smaller than that of the black hole but sufficiently large to perturb it. In this case, the generic stress-energy tensor $t_{\mu\nu}$ describing the matter-sources can be similarly decomposed in odd and even-parity parts

$$t_{\mu\nu} = \sum_{\ell, m} \left[(t_{\mu\nu}^{\ell m})^{(o)} + (t_{\mu\nu}^{\ell m})^{(e)} \right], \quad (10)$$

that are naturally gauge-invariant since the background is the vacuum Schwarzschild spacetime and are given explicitly by

$$(t_{\mu\nu}^{\ell m})^{(o)} = \begin{pmatrix} 0 & L_A^{\ell m} S_c^{\ell m} \\ L_A^{\ell m} S_c^{\ell m} & L^{\ell m} \nabla_{(d} S_{c)}^{\ell m} \end{pmatrix}, \quad (11)$$

for the odd-parity part and by

$$(t_{\mu\nu}^{\ell m})^{(e)} = \left(\begin{array}{c|c} T_{AB}^{\ell m} Y^{\ell m} & T_A^{\ell m} \nabla_c Y^{\ell m} \\ \hline T_A^{\ell m} \nabla_c Y^{\ell m} & r^2 T_3^{\ell m} Y^{\ell m} \gamma_{cd} + T_2^{\ell m} Z_{cd}^{\ell m} \end{array} \right), \quad (12)$$

for the even-parity one. Note that we have now used the Zerilli-Mathews set of harmonics for the expansion, that the ten coefficients $L_A^{\ell m}, L^{\ell m}, T_{AB}^{\ell m}, T_A^{\ell m}, T_2^{\ell m}, T_3^{\ell m}$ are gauge-invariant and that explicit expressions for them will be presented in the following Sections.

With the perturbations decomposed in a convenient form, we can now consider the Einstein field equations that, in the static vacuum background, take the simple form

$$\overset{\circ}{R}_{\mu\nu} = 0, \quad (13)$$

where $\overset{\circ}{R}$ is the Ricci tensor built from the background metric $\overset{\circ}{g}$. At first order in the perturbations, the linearity can be exploited to break up the Einstein equations as

$$\overset{\circ}{R}_{\mu\nu} + R_{\mu\nu} - \frac{1}{2} \overset{\circ}{g}_{\mu\nu} R = 8\pi t_{\mu\nu}, \quad (14)$$

where $\mathbf{R}$ is now the Ricci tensor built from the metric perturbations $\mathbf{h}$. Using equations (13), the field equations then reduce to

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}^0 R = 8\pi t_{\mu\nu} , \quad (15)$$

and where the components of the Ricci tensor are explicitly given by

$$R_{\mu\nu} = \nabla_\beta \Gamma_{\mu\nu}^\beta - \nabla_\nu \Gamma_{\mu\beta}^\beta . \quad (16)$$

Here, Γ are the Christoffel symbols relative to the perturbed metric $\mathbf{h}$, i.e.

$$\Gamma_{\mu\nu}^\beta = \frac{1}{2}g^{\alpha\beta}_0 (\partial_\nu h_{\mu\alpha} + \partial_\mu h_{\nu\alpha} - \partial_\alpha h_{\mu\nu}) , \quad (17)$$

and should be distinguished from the corresponding Christoffel symbols $\overset{0}{\Gamma}$ relative to the background metric $\overset{0}{g}$.

Note that while a generic perturbation will be a mixture of odd and even-parity contributions, we will exploit the linearity of the approach to handle them separately and simplify the treatment. In the following two Sections we will discuss the form the Einstein equations (15) assume in response to odd and even-parity perturbations over a Schwarzschild background. In particular, we will show how the three odd-parity coefficients of the expansion in harmonics of the metric and the seven even-parity ones can be combined to give two gauge-invariant master equations, named after Regge and Wheeler [4] and Zerilli [42], each of which is a wave-like equation in a scattering potential[‡].

Although our attention is here focussed on the *radiative* degrees of freedom of the perturbations (i.e. those with $\ell \geq 2$) because of their obvious application to the modelling of sources of gravitational waves, a comment should be made also on lower-order multipoles. In particular, it is worth remarking that the monopole component of the metric for a vacuum perturbation (i.e. with $\ell = 0$) is only of even-parity type and represents a variation in the mass-parameter of the Schwarzschild solution. On the other hand, the dipole component of the even-parity metric for a vacuum perturbation (i.e. with $\ell = 1$) is of pure-gauge type and it can be removed by means of a suitable gauge transformation [13]. Note that this is not the case for a dipolar odd-parity metric perturbation, which can instead be associated to the introduction of angular momentum onto the background metric. A detailed discussion of these low multipoles can be found in Appendix G of [13] and in [49].

4. Gauge-invariant odd-parity perturbations

Before discussing the derivation of the odd-parity equation a choice should be made for the odd-parity master function. Unfortunately, this choice has not been unique over the years and different authors have made different choices, making comparisons between different approaches less straightforward. Here, we will make a choice which highlights the relation with the gravitational-wave amplitude measured by a distant observer and, in particular, we construct the gauge-invariant combination of multipoles [19, 39, 50]

$$k_A \equiv h_A - \nabla_A h + 2h \frac{\nabla_A r}{r} , \quad (18)$$

[‡] These results were originally obtained by Regge and Wheeler [4] and by Zerilli [12, 13] in a specific gauge (i.e. the Regge-Wheeler gauge). Subsequently, the work of Moncrief showed how to reformulate the problem in a gauge-invariant form by deriving the equations from a suitable variational principle [14]. Some details of Moncrief's approach to odd-parity perturbations are briefly described in Appendix C.

where, we recall, ∇_A represents the covariant derivative with respect to the connection of the submanifold M^2 . In this way, the function

$$\Phi^{(o)}(t, r) \equiv r^3 \epsilon^{AB} \nabla_B \left(\frac{k_A}{r^2} \right) = r \left[\partial_t h_1^{(o)} - r^2 \partial_r \left(\frac{h_0^{(o)}}{r^2} \right) \right], \quad (19)$$

where ϵ_{AB} is the antisymmetric volume form on M^2 ($\epsilon^{01} = -\epsilon^{10} = -1$, see Appendix C), is gauge-invariant and will be our choice for the Regge-Wheeler master function [19, 39, 40, 50].

A slight variation of the master function (19) has been introduced by Cunningham, Price and Moncrief [15] in terms of the function $\tilde{\psi} \equiv \Lambda \Phi^{(o)}$ and this has been used so extensively in the literature [23, 24, 26] that it is now commonly referred to as the Cunningham-Price-Moncrief (CPM) convention. We partly follow this suggestion and introduce a new master function for the odd-parity perturbations defined as

$$\Psi^{(o)} \equiv \frac{1}{\Lambda - 2} \Phi^{(o)}. \quad (20)$$

With the choice (20), the Einstein field equations (15) with odd-parity perturbations lead to the inhomogeneous ‘Regge-Wheeler’ equation

$$\partial_t^2 \Psi^{(o)} - \partial_{r_*}^2 \Psi^{(o)} + V_\ell^{(o)} \Psi^{(o)} = S^{(o)}, \quad (21)$$

where

$$r_* \equiv r + 2M \ln \left(\frac{r}{2M} - 1 \right), \quad (22)$$

is the ‘tortoise coordinate’ [11] and $V_\ell^{(o)}$ is the odd-parity potential, defined as

$$V_\ell^{(o)} \equiv \left(1 - \frac{2M}{r} \right) \left(\frac{\Lambda}{r^2} - \frac{6M}{r^3} \right). \quad (23)$$

The right-hand-side of equation (21) represents the generic odd-parity ‘matter-source’ and is given by

$$S^{(o)} \equiv \frac{16\pi r}{\Lambda - 2} e^{2a} \epsilon^{AB} \nabla_B L_A = \frac{16\pi r}{\Lambda - 2} \left[\left(1 - \frac{2M}{r} \right) \partial_t L_1^{\ell m} - \partial_{r_*} L_0^{\ell m} \right], \quad (24)$$

with the components of the odd-parity matter-source vector defined as

$$L_A^{\ell m} \equiv \frac{1}{\Lambda} \int \frac{d\Omega}{\sin \theta} (i m t_{A2} Y_{\ell m}^* + t_{A3} \partial_\theta Y_{\ell m}^*), \quad A = 0, 1, \quad (25)$$

and where $d\Omega = \sin \theta d\phi d\theta$ is the surface element on the 2-sphere S^2 .

As a result of this construction, given a generic odd-parity matter-source $t_{\mu\nu}$ whose dynamics is known (*e.g.* through the solution of the equations of motion if the source is point-like, or those of relativistic hydrodynamics if the source is extended), equation (21) describes the evolution of the perturbations induced on the black-hole spacetime by the dynamics of the matter-source.

It should be noted that another choice for the gauge-invariant odd-parity master variable is possible and indeed was originally proposed by Moncrief [14]. This function, which hereafter we will refer to as the odd-parity Moncrief function, is defined as

$$Q^{(o)} \equiv \frac{\nabla_A r}{r} g^{AB} k_B = \frac{1}{r} \left(1 - \frac{2M}{r} \right) \left[h_1^{(o)} + \frac{r^2}{2} \partial_r \left(\frac{h_2}{r^2} \right) \right], \quad (26)$$

where the first expression is coordinate independent [49], while the second one is specialized to Schwarzschild coordinates with $h_2 = -2h$ [14]. In the Regge-Wheeler gauge $h_2 = h = 0$, and the definition (26) coincides with the variable used by Regge and Wheeler [4].

Historically, the choice of (26) as master variable has been the most common in the literature to describe odd-parity perturbations of a Schwarzschild spacetime and we will refer to it as ‘Regge-Wheeler’ (RW) convention. It should be noted that while (26) is a solution of the Regge-Wheeler equation, the corresponding source term differs from expression (24). A general expression of the source in the RW convention can be found in [49, 52] together with its specification for a point-particle (see also [53, 54, 55]).

The two master functions $Q^{(o)}$ and $\Psi^{(o)}$ are intimately related in at least two different ways. Firstly, through the variational formalism employed by Moncrief in [14] and a brief discussion of these relations is presented in Appendix C. Secondly, through an explicit expression that reads [49]

$$\partial_t \Psi^{(o)} = -Q^{(o)} + \frac{16\pi}{\Lambda - 2} \frac{r}{e^{2b}} L_1^{\ell m} . \quad (27)$$

Equation (27) highlights an important difference between the two master functions which is not just a dimensional one (i.e. $\Psi^{(o)}$ has the dimensions of a length, while $Q^{(o)}$ is dimensionless) and that will have consequences on the asymptotic expressions for the gravitational waveforms when these are expressed in one or in the other convention. A detailed discussion of this will be made in sections. 6.1, 6.3 and 6.4.

5. Gauge-invariant even-parity perturbations

Also in the case of even-parity perturbations, an evolution equation similar to the Regge-Wheeler one (21) can be found. In particular, following Moncrief [14], we define the functions

$$\kappa_1 \equiv K + \frac{1}{e^{2b}} \left(r \partial_r G - \frac{2}{r} h_1^{(e)} \right) , \quad (28)$$

$$\kappa_2 \equiv \frac{1}{2} [e^{2b} H_2 - e^b \partial_r (r e^b K)] , \quad (29)$$

where both κ_1 and κ_2 are gauge-invariant functions, as well as the following linear combination

$$q_1 \equiv r \Lambda \kappa_1 + \frac{4r}{e^{4b}} \kappa_2 . \quad (30)$$

Strictly related to expression (30) is the gauge-invariant function most frequently used in the literature [25, 41, 52, 56, 57, 58, 59, 60, 62]§

$$\Psi^{(e)} \equiv \frac{r q_1}{\Lambda [r (\Lambda - 2) + 6M]} . \quad (31)$$

which is also the solution of the inhomogeneous even-parity master equation or ‘Zerilli’ equation

$$\partial_t^2 \Psi^{(e)} - \partial_{r_*}^2 \Psi^{(e)} + V_\ell^{(e)} \Psi^{(e)} = S^{(e)} , \quad (32)$$

and, again, is a wave-like equation in the scattering Zerilli potential [12]

$$V_\ell^{(e)} \equiv \left(1 - \frac{2M}{r} \right) \frac{\Lambda(\Lambda - 2)^2 r^3 + 6(\Lambda - 2)^2 M r^2 + 36(\Lambda - 2) M^2 r + 72 M^3}{r^3 [(\Lambda - 2) r + 6M]^2} . \quad (33)$$

§ Note that expression (31) corrects the sign in the corresponding definition of the even-parity function used in [63] (cf equation(7) in [63]).

The even-parity matter-source has a rather extended expression given by

$$S^{(e)} = -\frac{8\pi}{\Lambda[(\Lambda-2)r+6M]} \left\{ \frac{\Lambda(6r^3-16Mr^2)-r^3\Lambda^2-8r^3+68Mr^2-108M^2r}{(\Lambda-2)r+6M} T_{00}^{\ell m} \right. \\ \left. + \frac{1}{e^{4b}} [2Mr+r^2(\Lambda-4)] T_{11}^{\ell m} + 2r^3 \partial_{r_*} T_{00}^{\ell m} - \frac{2r^3}{e^{4b}} \partial_{r_*} T_{11}^{\ell m} \right. \\ \left. + \frac{4\Lambda r}{e^{4b}} T_1^{\ell m} + \frac{1}{e^{2b}} \left[2\Lambda \left(1 - \frac{3M}{r} \right) - \Lambda^2 \right] T_2^{\ell m} + \frac{4r^2}{e^{4b}} T_3^{\ell m} \right\}. \quad (34)$$

and we refer the interested reader to [52, 60, 61] where the steps necessary for the derivation of equation (34) are given in detail and some application to accretion problems are discussed. Here, we simply recall that the expressions of the even-parity vector and tensor spherical-harmonics for the matter-source needed in (34) can be obtained from the orthogonality properties of the harmonics and are

$$T_A^{\ell m} = \frac{1}{\Lambda} \int d\Omega \left[t_{A2} \partial_\theta (Y_{\ell m}^*) - t_{A3} \frac{im Y_{\ell m}^*}{\sin^2 \theta} \right], \quad A = 0, 1, \quad (35)$$

$$T_2^{\ell m} = \frac{2}{\Lambda(\Lambda-2)} \int d\Omega \left[t_{22} \frac{W_{\ell m}^*}{2} + t_{23} \frac{2X_{\ell m}^*}{\sin \theta} + t_{33} \left(\frac{\Lambda Y_{\ell m}^*}{2} - \frac{m^2 Y_{\ell m}^*}{\sin^2 \theta} + \cot \theta \partial_\theta Y_{\ell m}^* \right) \right], \quad (36)$$

$$T_3^{\ell m} = \frac{1}{2r^2} \int d\Omega \left(t_{22} + t_{33} \frac{1}{\sin^2 \theta} \right) Y_{\ell m}^*, \quad (37)$$

$$T_{AB}^{\ell m} = \int d\Omega t_{AB} Y_{\ell m}^*, \quad A, B = 0, 1, \quad (38)$$

where the angular functions $W^{\ell m}(\theta, \phi)$ and $X^{\ell m}(\theta, \phi)$ are defined as [4]

$$W^{\ell m} \equiv \frac{\nabla_{(\phi} S_{\theta)}^{\ell m}}{\sin \theta} = \partial_\theta^2 Y^{\ell m} - \cot \theta \partial_\theta Y^{\ell m} - \frac{1}{\sin^2 \theta} \partial_\phi^2 Y^{\ell m}, \quad (39)$$

$$X^{\ell m} \equiv -\sin \theta \left(\nabla_\theta S_\theta^{\ell m} - \frac{\nabla_\phi S_\phi^{\ell m}}{\sin^2 \theta} \right) = 2 \left(\partial_{\theta\phi}^2 Y^{\ell m} - \cot \theta \partial_\phi Y^{\ell m} \right). \quad (40)$$

and where the asterisk stands for complex conjugation. A few comments are now needed about the different notations in which the even-parity functions can be found in the literature. A particularly common choice is that proposed by Moncrief in [14] for the the even-parity gauge-invariant master function $Q^{(e)}$ which is related to Zerilli function (31) simply as $Q^{(e)} = \Lambda \Psi^{(e)}$. Other authors (e.g. [52, 56]) use instead a master function defined as $Z \equiv 2\Psi^{(e)}$ and a factor of 2 needs to be introduced or removed when converting between the two notations. Yet another even-parity function can be introduced in terms of two new gauge-invariant metric functions k and χ where [40, 41]

$$k = \kappa_1 = K + \frac{1}{e^{2b}} \left[r \partial_r G - \frac{2}{r} h_1^{(e)} \right], \quad (41)$$

$$\chi + k = H_2 - \frac{2}{e^{2b}} \partial_r h_1^{(e)} - \frac{2M}{r^2} h_1^{(e)} + \frac{1}{e^{2b}} \partial_r (r^2 \partial_r G) + M \partial_r G, \quad (42)$$

and such that

$$\kappa_2 = \frac{1}{2} e^{2b} \left(\chi - r \partial_r k + \frac{M}{r} e^{2b} k \right). \quad (43)$$

In this case, the Zerilli function (31) can be equivalently defined as [41]

$$\Psi^{(e)} \equiv \frac{2r^2}{\Lambda[(\Lambda-2)r+6M]e^{2b}} \left[\chi + \left(\frac{\Lambda}{2} + \frac{M}{r} \right) e^{2b} k - r \partial_r k \right]. \quad (44)$$

Finally, we recall that the homogeneous odd and even-parity master equations (21) and (32) can be transformed into each other by means of differential operations [1] and they are connected to the master equation that Bardeen and Press [65] have derived via the Newman-Penrose formalism.

5.1. Initial data

An important aspect in the solution in the time-domain of the perturbation equations (21) and (32) is the specification of the initial data, namely the determination of the Regge-Wheeler and Zerilli functions $\Psi^{(o)}$ and $\Psi^{(e)}$ at a given initial time and in the presence of the external sources of perturbations. The standard approach in this case is that of exploiting the $1+1$ splitting of spacetime and to solve the perturbed Hamiltonian and momentum-constraint equations as a set of coupled ordinary differential equations having the matter perturbations as sources.

In the case of even-parity perturbations, in particular, the Hamiltonian constraint takes the form [39, 40]

$$\partial_{r_*}^2 k + \frac{2-5M}{r} \partial_{r_*} k - \frac{r-2M}{r^2} \partial_{r_*} \chi - \frac{r-2M}{r^3} \left[\Lambda k - \left(\frac{\Lambda}{2} + 1 \right) \chi \right] = -8\pi T_{00}^{\ell m}, \quad (45)$$

and the momentum constraint is instead given by

$$\begin{aligned} \partial_{r_*}^2 \dot{k} + \frac{2(r-2M)}{r^2} \partial_{r_*} \dot{k} - \frac{2M^2 + r\Lambda(r-2M)}{2r^4} \dot{k} + \left(1 - \frac{2M}{r} \right) \frac{4(r-2M) - \Lambda r}{4r^3} \dot{\chi} \\ - \frac{r-2M}{r^2} \partial_{r_*} \dot{\chi} = \frac{4\pi}{r-2M} \left\{ 4r \partial_{r_*} T_{01}^{\ell m} + \frac{1}{r} [\Lambda T_0^{\ell m} + 4(2r-3M) T_{01}^{\ell m}] \right\}, \quad (46) \end{aligned}$$

where $\dot{k} \equiv \partial_t k$ and $\dot{\chi} \equiv \partial_t \chi$.

Clearly, equations (45) and (46) are not sufficient to determine all of the unknowns: i.e. $(k, \dot{k})$ and $(\chi, \dot{\chi})$ and this degree of indeterminacy reflects the arbitrary amount of gravitational radiation that can be present at the initial time with the same matter-sources. To alleviate at least in part this problem, the initial data is chosen so as to be best adapted to the physical problem under consideration. To minimize the initial gravitational-wave content, for instance, it is possible to set $\chi = 0 = \dot{\chi}$ (conformally-flat approximation) and to use eqs. (45) and (46) to compute k and $\dot{k}$. Alternatively, it is possible to consider time-symmetric initial data for which $\dot{k} = 0 = \dot{\chi}$ and compute χ and k from the constraint equations [59]. Finally, it is possible to specify a solution for χ and $\dot{\chi}$ and use the constrain equations to compute k and $\dot{k}$ [58].

Similarly, in the case of odd-parity perturbations, we can exploit the linearized Einstein equations [cf equations (18) and (20) of [39]]

$$g^{\circ CD} \nabla_D [4r(\nabla_{[A} r)k_{C]} + 2r^2 \nabla_{[C} k_{A]}] - (\Lambda - 2)k_A = -16\pi L_A^{\ell m}, \quad (47)$$

$$\nabla_A k^A = 16\pi L^{\ell m}, \quad (48)$$

where

$$L^{\ell m} \equiv \frac{1}{\Lambda(\Lambda-1)} \int \frac{d\Omega}{\sin \theta} \left[t_{23} W_{\ell m}^* + \frac{1}{2} \left(-t_{22} + \frac{1}{\sin^2 \theta} t_{33} \right) X_{\ell m}^* \right]. \quad (49)$$

Tedious but straightforward algebra allows to select two of the three equations (47) and (48) as constraint equations for the four unknowns $(k_0, \dot{k}_0)$ and $(k_1, \dot{k}_1)$

$$\partial_{r_*}^2 k_1 + \frac{2M}{r^2} \partial_{r_*} k_1 - \frac{4M}{r^2 e^{4b}} k_1 - e^{2b} \partial_{r_*} \dot{k}_0 + \frac{2M}{r^2} e^{2b} \dot{k}_0 = 16\pi \partial_{r_*} L^{\ell m} \quad (50)$$

$$\begin{aligned} \partial_{r_*}^2 k_0 + \frac{2(r-3M)}{r^2} \partial_{r_*} k_0 + \left(\frac{2M}{r^3} - \frac{\Lambda}{r^2} \right) \frac{1}{e^{2b}} k_0 + \frac{2M}{r^3 e^{4b}} k_1 - \frac{2}{r e^{2b}} \dot{k}_0 \\ - \frac{2}{r e^{4b}} \dot{k}_1 - \frac{1}{e^{2b}} \partial_{r_*} \dot{k}_1 = -\frac{16\pi}{e^{2b}} L_0^{\ell m}, \end{aligned} \quad (51)$$

where $\dot{k}_A \equiv \partial_t k_A$. As for the even-parity perturbations, the constraint equations (50) and (51) can be solved as a set of coupled ordinary differential equations to compute two of the four unknowns when the other two are specified according to the problem under consideration.

As a final remark we recall that an equivalent and alternative approach to the specification of the initial data consist in setting $\partial_t \Psi^{(o)} = 0 = \partial_t \Psi^{(e)}$ and in solving the Regge-Wheeler and Zerilli equations as ordinary differential equations [15, 16]. In the case of even-parity perturbations this is equivalent to setting $\dot{k} = 0 = \dot{\chi}$ (cf equation (44)) and thus considering time-symmetric initial data.

5.2. Frequency domain and QNMs

We conclude this Section by recalling that the investigation of the Regge-Wheeler and Zerilli equations (21)–(32) is often made in the frequency domain, where the properties of Fourier transform can be exploited and the oscillation properties of the spacetime are easier to interpret. What is usually done in this case is to assume a harmonic time dependence in the perturbations, i.e. $\Psi^{(e)} \sim \exp(i\omega_n t)$ and $\Psi^{(o)} \sim \exp(i\omega_n t)$ where ω_n is the complex oscillation frequency of the n -th mode, so that the Regge-Wheeler and Zerilli equations (21)–(32) take respectively the form

$$\left(\Psi^{(o)} \right)'' + \left(\omega_n^2 - V_\ell^{(o)} \right) \Psi^{(o)} = S^{(o)}, \quad (52)$$

$$\left(\Psi^{(e)} \right)'' + \left(\omega_n^2 - V_\ell^{(e)} \right) \Psi^{(e)} = S^{(e)}, \quad (53)$$

where a ‘prime’ indicates a total derivative with respect to r_* . Equations (52)–(53) can be solved as an eigenvalue problem with boundary conditions such that the perturbations behave as *pure outgoing-waves* at spatial infinity

$$\Psi^{(o)}, \Psi^{(e)} \sim \exp(-i\omega_n r_*) \quad \text{for } r_* \rightarrow +\infty. \quad (54)$$

and as *pure ingoing-waves* at the event horizon

$$\Psi^{(o)}, \Psi^{(e)} \sim \exp(i\omega_n r_*) \quad \text{for } r_* \rightarrow -\infty, \quad (55)$$

In the absence of source terms, the modes solutions of equations (52) and (53) are referred to as the quasi-normal modes (QNMs) of the black hole, with the real part of ω_n representing the oscillation frequency and with the imaginary part of ω_n^{-1} representing the damping time of the oscillation. A complete discussion of the properties and astrophysical relevance of these modes can be found in [1, 2, 3] and here we only recall the most important properties for a Schwarzschild black hole:

- All of the QNMs have positive imaginary parts and represent therefore damped modes. Stated differently, a Schwarzschild black hole is *linearly stable* against perturbations.

- The damping time of QNMs depends linearly on the mass of the black hole and is shorter for higher-order modes.
- The excitation of a black hole is referred to as its “ringing” and the late-time dynamics of these perturbations (i.e. the ‘tail’ of the ringing) can be described with a power-law representing the envelope of the various decaying QNMs.
- The QNMs are *isospectral*, i.e. odd and even-parity perturbations have the same complex eigenfrequencies. This is due to the uniqueness in which a black hole can react to a perturbation and it is not true for a relativistic star.

6. Asymptotic expressions and gravitational waves

In the previous Section we have reviewed the derivation of the equations describing the evolution of perturbations of nonrotating black holes induced by a non-zero stress-energy tensor. These perturbations have been assumed to be generic in nature, needing to satisfy only the condition of having a mass-energy much smaller than that of the black hole. The solution of these equations with suitable initial conditions completely specifies the reaction of the black hole to the perturbations and this is essentially represented by the emission of gravitational waves.

As mentioned in Section 2, the importance of the gauge-invariant variables used so far is that they are directly related to the amplitude and energy of the gravitational-wave signal measured at large distances. The purpose of this Section is to review the steps necessary to obtain the relations between the master functions for the odd and even-parity perturbations and the ‘plus’ and ‘cross’ polarization amplitudes h_+ , $h_\times$ of a gravitational wave in a transverse and traceless (TT) gauge. In practice, and following the guidelines tracked in [15, 16], we will derive an expression for the perturbation metric h equivalent to that obtained in the standard TT-gauge on a Minkowski spacetime and relate it to the odd and even-parity master functions $\Psi^{(o)}$ and $\Psi^{(e)}$.

To obtain this result a number of conditions need to be met. Firstly, we need to evaluate each multipole of the decomposed metric perturbations in the tetrad e of stationary observers in the background Schwarzschild spacetime, i.e. $h_{\hat{\mu}\hat{\nu}} = e_{\hat{\mu}}^\mu e_{\hat{\nu}}^\nu h_{\mu\nu}$, where e is diagonal with components $e_{\hat{\mu}}^\mu \equiv \{e^b, e^{-b}, r^{-1}, (r \sin \theta)^{-1}\}$ and where the indices $\hat{\mu}$ refer to the locally ‘flat’ coordinates. Secondly, all of the quantities need to be evaluated far away from the source (i.e. in the “wave-zone”) and in the so-called *radiation gauge*. In practice, this amounts to requiring that components $h_{\hat{\theta}\hat{\theta}}$, $h_{\hat{\phi}\hat{\phi}}$ and $h_{\hat{\theta}\hat{\phi}}$ are functions of the type $f(t - r)/r$ (i.e. they are outgoing spherical waves), while all the other components have a more rapid decay of $\mathcal{O}(1/r^2)$. Thirdly, we need to impose the condition that the metric is traceless modulo higher order terms, i.e. $h_{\hat{\theta}\hat{\theta}} + h_{\hat{\phi}\hat{\phi}} = 0 + \mathcal{O}(1/r^2)$.

6.1. Asymptotic expressions from odd-parity perturbations

We first consider odd-parity perturbations and recall that from the radiation-gauge conditions and since for large r the metric asymptotes that of a flat spacetime $e^b \sim e^{-b} \sim 1$, we have

$$h_{\hat{\theta}\hat{t}}^{(o)} = \frac{h_0^{(o)}}{r} e^b S_\theta \sim \frac{h_0^{(o)}}{r} \sim \mathcal{O}\left(\frac{1}{r^2}\right) \longrightarrow h_0^{(o)} \sim \mathcal{O}\left(\frac{1}{r}\right), \quad (56)$$

$$h_{\hat{\theta}\hat{r}}^{(o)} = \frac{h_1^{(o)}}{r} e^b S_\theta \sim \frac{h_1^{(o)}}{r} \sim \mathcal{O}\left(\frac{1}{r^2}\right) \longrightarrow h_1^{(o)} \sim \mathcal{O}\left(\frac{1}{r}\right), \quad (57)$$

where the ℓ, m indices have been omitted for clarity. Similarly, since $h_{\hat{\theta}\hat{\theta}}^{(o)} = 2hr^{-2}\nabla_\theta S_\theta \sim \mathcal{O}(1/r)$ we deduce that $h \sim \mathcal{O}(r)$ so that the only components of the metric having wave-like properties at large r are

$$h_+^{(o)} \equiv \frac{1}{2} \left(h_{\hat{\theta}\hat{\theta}}^{(o)} - h_{\hat{\phi}\hat{\phi}}^{(o)} \right) = \frac{h}{r^2} \left(\nabla_\theta S_\theta - \frac{\nabla_\phi S_\phi}{\sin^2 \theta} \right) + \mathcal{O}\left(\frac{1}{r^2}\right), \quad (58)$$

$$h_\times^{(o)} \equiv h_{\hat{\theta}\hat{\phi}}^{(o)} = \frac{h}{r^2} \frac{\nabla_{(\phi} S_{\theta)}}{\sin \theta} + \mathcal{O}\left(\frac{1}{r^2}\right), \quad (59)$$

and where we note that since h has the dimensions of a length squared, both h_+ and $h_\times$ are, as expected, dimensionless.

Next, we need to relate the perturbation h to the odd-parity master function $\Psi^{(o)}$. To do so, we follow the procedure outlined in [15], and note that (cf equation (III-20) ||)

$$\partial_t h = \left(1 - \frac{2M}{r}\right) \partial_r \left(r\Psi^{(o)}\right) + h_0^{(o)}, \quad (60)$$

Equation (60) represents one of the Hamilton equations as derived by Moncrief in a Hamiltonian formulation of perturbation equations [14] (see Appendix C for details). We also note that the radiation-gauge conditions on h and $h_0^{(o)}$ imply that $\Psi^{(o)} \sim \mathcal{O}(1)$, i.e., in the wave-zone $\Psi^{(o)}$ has the dimensions of a length, behaves as an outgoing-wave, but it does not depend explicitly on r .

Exploiting now the outgoing-wave behaviour of h at large distances we can write

$$\partial_t h = -\partial_r h + \mathcal{O}\left(\frac{1}{r}\right), \quad (61)$$

so that asymptotically equation (60) simply becomes

$$\partial_r h = -\partial_r \left(r\Psi^{(o)}\right) + \mathcal{O}\left(\frac{1}{r}\right), \quad (62)$$

and its integration yields

$$\frac{h}{r} \sim -\Psi^{(o)} + \mathcal{O}\left(\frac{1}{r}\right). \quad (63)$$

As a result, the ‘+’ and ‘ $\times$ ’ polarization amplitudes of the gravitational wave can be calculated from eqs. (59)–(58) as

$$h_+^{(o)} = -\frac{1}{r}\Psi^{(o)} \left(\nabla_\theta S_\theta - \frac{\nabla_\phi S_\phi}{\sin^2 \theta} \right) + \mathcal{O}\left(\frac{1}{r^2}\right), \quad (64)$$

$$h_\times^{(o)} = -\frac{1}{r}\Psi^{(o)} \frac{\nabla_{(\phi} S_{\theta)}}{\sin \theta} + \mathcal{O}\left(\frac{1}{r^2}\right). \quad (65)$$

It is now interesting to note that expressions (64) and (65) can be written in a compact form using the $s = -2$ spin-weighted spherical harmonics

$$_{-2}Y^{\ell m}(\theta, \phi) \equiv \sqrt{\frac{(\ell-2)!}{(\ell+2)!}} \left(W^{\ell m} - i \frac{X^{\ell m}}{\sin \theta} \right), \quad (66)$$

so that expressions (64) and (65) can be combined into a single complex expression given by

$$\left(h_+^{(o)} - i h_\times^{(o)} \right)_{\ell m} = \frac{i}{r} \sqrt{\frac{(\ell+2)!}{(\ell-2)!}} \Psi_{\ell m}^{(o)} {}_{-2}Y^{\ell m}(\theta, \phi) + \mathcal{O}\left(\frac{1}{r^2}\right), \quad (67)$$

|| We recall that in the notation of [15] $\tilde{\psi} = \Lambda(\Lambda-2)\Psi^{(o)}$, and the multipoles in [15] are related to ours as $\tilde{h}_2 = h_2 = -2h$, $\tilde{h}_0 = h_0^{(o)}$ and $\tilde{h}_1 = h_1^{(o)}$

where, for clarity, we have explicitly restored the multipole indices ℓ, m .

An important clarification is now needed. We have discussed in the previous Section that the odd-parity metric perturbations are sometimes expressed in terms of the odd-parity Moncrief function $Q^{(o)}$ (cf equation (26)) and it is not unusual to find in the literature the gravitational wave amplitudes expressed in terms of this quantity. However, great care must be paid to the asymptotic relation between the master function $Q^{(o)}$ and the gravitational-wave amplitudes and, indeed, this is sometimes source of confusion [63, 64]. To clarify this point, we recall that the derivation of the asymptotic relation between $Q^{(o)}$ and h proceeds as in a way similar to the one discussed above. In the radiation gauge and at large distances from the black hole, we can use relation (61) in the definition (26) with $h_2 = -2h$, so that

$$Q^{(o)} \sim \frac{1}{r} \partial_t h + \mathcal{O}\left(\frac{1}{r}\right), \quad (68)$$

which is also a dimensionless quantity. Since $h \sim \mathcal{O}(r)$, the function $Q^{(o)}$ does not depend on r at leading order and equation (68) can be integrated to give

$$\frac{h(t)}{r} \sim \int_{-\infty}^t Q^{(o)}(t') dt' + \mathcal{O}\left(\frac{1}{r}\right) + \text{const.}, \quad (69)$$

where the integration constant can be defined as

$$\text{const.} \equiv \lim_{t \rightarrow -\infty} \frac{h(t, r)}{r} \sim \mathcal{O}(1), \quad (70)$$

and it can be set to zero in the case of asymptotically flat metric perturbations ($h = 0$) at earlier times. Combining now expressions (39), (40) and (69), the gravitational-wave amplitudes in the two polarizations and with the new master function read

$$\left(h_+^{(o)} - i h_\times^{(o)}\right)_{\ell m} = -\frac{i}{r} \sqrt{\frac{(\ell+2)!}{(\ell-2)!}} \left(\int_{-\infty}^t Q_{\ell m}^{(o)}(t') dt'\right) {}_{-2}Y^{\ell m}(\theta, \phi) + \mathcal{O}\left(\frac{1}{r^2}\right). \quad (71)$$

Comparing expressions (67) and (71) we note that while $\Psi^{(o)}$ and $Q^{(o)}$ are both solutions of the Regge-Wheeler equation (21), they yield two different asymptotic expressions for the gravitational-wave amplitudes. This difference, which is consistent with equation (27) when evaluated in an asymptotic region of the spacetime where $L_1^{\ell m} = 0$, is subtle but important and, as mentioned above, it has led to some inconsistencies in the literature both for the determination of the asymptotic gravitational-wave amplitudes and for the energy losses. This will be further discussed in sections 6.3 and 6.4.

6.2. Asymptotic expressions from even-parity perturbations

A calculation conceptually analogous to the one carried out in sections 6.1 leads to the relation between the gravitational-wave amplitude and the even-parity master function. In particular, after projecting equation (8) along the stationary tetrad, the asymptotic wave amplitudes in the two polarization states are

$$h_+^{(e)} = \frac{1}{2} \left(h_{\hat{\theta}\hat{\theta}}^{(e)} - h_{\hat{\phi}\hat{\phi}}^{(e)}\right) = \frac{G}{2} \left(\nabla_\theta \nabla_\theta Y^{\ell m} - \frac{\nabla_\phi \nabla_\phi Y^{\ell m}}{\sin^2 \theta}\right) = \frac{G}{2} W^{\ell m}, \quad (72)$$

$$h_\times^{(e)} = h_{\hat{\theta}\hat{\phi}}^{(e)} = G \frac{\nabla_\theta \nabla_\phi Y^{\ell m}}{\sin \theta} = \frac{G}{2} \frac{X^{\ell m}}{\sin \theta}, \quad (73)$$

so that we essentially need to relate the metric perturbation G with the even-parity function $\Psi^{(e)}$. Firstly, it is easy to realize that the even-parity metric projected onto the tetrad, $h_{\hat{\mu}\hat{\nu}}^{(e)}$ is such that

$$H_2 \sim \mathcal{O}\left(\frac{1}{r^2}\right), \quad \text{and} \quad h_1^{(e)} \sim \mathcal{O}\left(\frac{1}{r}\right), \quad (74)$$

so that the terms proportional to these multipoles are of higher order for large r and can be neglected. Furthermore, from the traceless condition

$$h_{\hat{\theta}\hat{\theta}}^{(e)} + h_{\hat{\phi}\hat{\phi}}^{(e)} = 0 + \mathcal{O}\left(\frac{1}{r^2}\right), \quad (75)$$

we obtain an asymptotic relation between K and G of the type

$$2KY^{\ell m} + G\left(\nabla_{\theta}\nabla_{\theta}Y^{\ell m} + \frac{\nabla_{\phi}\nabla_{\phi}Y^{\ell m}}{\sin^2\theta}\right) = (2K - G\Lambda)Y^{\ell m} \sim \mathcal{O}\left(\frac{1}{r^2}\right), \quad (76)$$

where we have used the definition (5) to derive the right-hand-side of expression (76). As a result, the asymptotic relation between the two components of the even-parity part of the perturbation metric simply reads

$$K \sim \frac{\Lambda}{2}G + \mathcal{O}\left(\frac{1}{r^2}\right). \quad (77)$$

Using now the definitions (28)–(29), we have that asymptotically

$$\kappa_1 \sim \frac{\Lambda}{2}G + r\partial_r G + \mathcal{O}\left(\frac{1}{r^2}\right) \quad (78)$$

$$\kappa_2 \sim -\frac{1}{2}(K + r\partial_r K) \sim -\frac{\Lambda}{4}(G + r\partial_r G) + \mathcal{O}\left(\frac{1}{r^2}\right), \quad (79)$$

and their linear combination (29) becomes

$$q_1 \sim \frac{rG}{2}\Lambda(\Lambda - 2) + \mathcal{O}\left(\frac{1}{r}\right). \quad (80)$$

Finally, the asymptotic gauge-invariant even-parity master function reads

$$\Psi^{(e)} \sim \frac{rq_1}{\Lambda[r(\Lambda - 2) + 6M]} \sim \frac{1}{2}rG + \mathcal{O}\left(\frac{1}{r}\right), \quad (81)$$

so that, modulo higher-order terms, the even-parity gravitational-wave amplitudes measured by a distant observer can be written in the compact form

$$\left(h_+^{(e)} - ih_{\times}^{(e)}\right)_{\ell m} = \frac{1}{r}\sqrt{\frac{(\ell - 2)!}{(\ell + 2)!}}\Psi_{\ell m - 2}^{(e)}Y^{\ell m}(\theta, \phi) + \mathcal{O}\left(\frac{1}{r^2}\right). \quad (82)$$

6.3. Asymptotic general expressions

It is often convenient to combine the expressions for the asymptotic gravitational wave amplitudes due to odd and even-parity perturbations into a single expression of the type

$$h_+ - ih_{\times} = \frac{1}{r}\sum_{\ell, m}\sqrt{\frac{(\ell + 2)!}{(\ell - 2)!}}\left(\Psi_{\ell m}^{(e)} + i\Psi_{\ell m}^{(o)}\right)_{-2}Y^{\ell m}(\theta, \phi) + \mathcal{O}\left(\frac{1}{r^2}\right), \quad (83)$$

or, equivalently

$$h_+ - ih_\times = \frac{1}{r} \sum_{\ell, m} \sqrt{\frac{(\ell+2)!}{(\ell-2)!}} \left(\Psi_{\ell m}^{(e)} - i \int_{-\infty}^t Q_{\ell m}^{(o)}(t') dt' \right) {}_{-2}Y^{\ell m}(\theta, \phi) + \mathcal{O}\left(\frac{1}{r^2}\right), \quad (84)$$

where we have defined $h_+ \equiv h_+^{(o)} + h_+^{(e)}$ and $h_\times \equiv h_\times^{(o)} + h_\times^{(e)}$. Note that $X^{\ell 0} = 0$ for any value of ℓ , so that in the case of axisymmetry the gravitational wave signal is proportional to $W^{\ell 0}$ only.

It is also useful to underline that while expression (83) resembles the corresponding expression (10) of [63], it is indeed very different. Firstly, because in [63] the Moncrief function is adopted for the odd-parity part of the perturbations and hence, modulo a normalisation factor, the function $\Psi^{(o)}$ appearing there corresponds to our function $Q^{(o)}$ (cf expression (26)). Secondly, because with this choice for the odd-parity perturbations a time derivative is needed in the asymptotic expression for the gravitational-wave amplitudes (cf the discussion in the derivation of equation (71)). As a result, expression (10) of [63] (which is also missing the distinction between the real and imaginary parts) should really be replaced by our expression (84). A similar use of the Moncrief function for the odd-parity part is present also in [66, 67, 68], where it is employed to calculate the gravitational-wave content of numerically simulated spacetimes.

6.4. Energy and angular momentum losses

Using the expressions derived so far we can now estimate the energy and angular momentum losses due to gravitational waves propagating outwards to spatial infinity. More specifically, this can be done by using expression (83) and the definition of Isaacson's stress-energy pseudo-tensor $\tau_{\mu\nu}$ for the gravitational-wave field h propagating in the curved background field $\bar{g}$ and in a Lorentz gauge [69, 70]

$$\tau_{\mu\nu} \equiv \frac{1}{32\pi} \langle \nabla_\mu h_{\alpha\beta} \nabla_\nu h^{\alpha\beta} \rangle, \quad (85)$$

where the brackets $\langle \dots \rangle$ refer to a spatial average over a length much larger than the typical gravitational wavelength [69, 11]. The averaged expression (85) is gauge-invariant [69] and holds in the 'limit of high frequency' (or *short-wave* approximation), i.e. whenever the wavelength of the gravitational-wave field is small when compared to the local radius of curvature of the background geometry. In practice, gravitational radiation from isolated systems is of high frequency whenever it is far enough away from its source.

Expression (85) accounts for the amount of energy and momentum carried by the gravitational wave over a certain region of spacetime, but since we are interested in the energy flux as measured by an inertial observer, we need to project the pseudo-tensor on the observer's locally orthonormal tetrad, where it becomes

$$\tau_{\hat{\mu}\hat{\nu}} \equiv \frac{1}{32\pi} \langle \partial_{\hat{\mu}} \bar{h}_{\hat{\alpha}\hat{\beta}} \partial_{\hat{\nu}} \bar{h}^{\hat{\alpha}\hat{\beta}} \rangle, \quad (86)$$

with $\bar{h}_{\hat{\mu}\hat{\nu}} \equiv h_{\hat{\mu}\hat{\nu}} - \frac{1}{2} \eta_{\hat{\mu}\hat{\nu}}$. As a result, the energy per unit time and angle carried by the gravitational waves and measured by a stationary observer at large distance from the black hole is given by

$$\frac{d^2 E}{dt d\Omega} = \frac{r^2}{16\pi} \left[\left(\frac{dh_{\hat{\theta}\hat{\phi}}}{dt} \right)^2 + \frac{1}{4} \left(\frac{dh_{\hat{\theta}\hat{\theta}}}{dt} - \frac{dh_{\hat{\phi}\hat{\phi}}}{dt} \right)^2 \right] = \frac{r^2}{16\pi} \left(\left| \frac{dh_+}{dt} \right|^2 + \left| \frac{dh_\times}{dt} \right|^2 \right), \quad (87)$$

where the total derivative is made with respect to the asymptotic observer's time. Integrating (87) over the solid angle, the total power emitted in gravitational waves is then given by

$$\frac{dE}{dt} = \frac{1}{16\pi} \sum_{\ell,m} \frac{(\ell+2)!}{(\ell-2)!} \left(\left| \frac{d\Psi_{\ell m}^{(e)}}{dt} \right|^2 + \left| \frac{d\Psi_{\ell m}^{(o)}}{dt} \right|^2 \right), \quad (88)$$

$$= \frac{1}{16\pi} \sum_{\ell,m} \left(\Lambda(\Lambda-2) \left| \frac{d\Psi_{\ell m}^{(e)}}{dt} \right|^2 + \frac{\Lambda}{\Lambda-2} \left| \frac{d\Psi_{\ell m}^{(o)}}{dt} \right|^2 \right), \quad (89)$$

where expression (89) was first presented in [15, 16].

For the same reasons discussed in the previous section, these expressions need to be suitably modified when the energy losses are expressed in terms of the odd-parity Moncrief function $Q^{(o)}$, in which case they assume the form

$$\frac{dE}{dt} = \frac{1}{16\pi} \sum_{\ell,m} \frac{(\ell+2)!}{(\ell-2)!} \left(\left| \frac{d\Psi_{\ell m}^{(e)}}{dt} \right|^2 + \left| Q_{\ell m}^{(o)} \right|^2 \right). \quad (90)$$

Besides having arbitrary angular dependence, the perturbations can also have non-diagonal terms and thus account for a non-zero net angular momentum, part of which can be lost to gravitational radiation. In this case, the angular momentum flux carried away in the form of gravitational waves can be calculated in terms of the $r\phi$ component of the stress energy tensor (86). In particular, using spherical coordinates and assuming that the rotation is parametrised by the angle ϕ , we have

$$\frac{d^2 J}{dt d\Omega} = \frac{r^2}{32\pi} \langle \partial_\phi \bar{h}_{\hat{\mu}\hat{\nu}} \partial_r \bar{h}^{\hat{\mu}\hat{\nu}} \rangle = -\frac{r^2}{16\pi} \langle \partial_r h_{\hat{\theta}\hat{\theta}} \partial_\phi h_{\hat{\theta}\hat{\theta}} + \partial_r h_{\hat{\theta}\hat{\phi}} \partial_\phi h_{\hat{\theta}\hat{\phi}} \rangle. \quad (91)$$

Since the metric components with the radiation-gauge condition behave like outgoing spherical waves and since $h_{\hat{\theta}\hat{\theta}} = h_+$ and $h_{\hat{\theta}\hat{\phi}} = h_\times$, the angular momentum carried away in the form of gravitational waves (91) is then expressed as

$$\frac{d^2 J}{dt d\Omega} = -\frac{r^2}{16\pi} (\partial_t h_+ \partial_\phi h_+^* + \partial_t h_\times \partial_\phi h_\times^*), \quad (92)$$

Proceeding in a way similar to the one followed in the calculation of the emitted power, the total angular momentum lost per unit time to gravitational wave reads [49]

$$\frac{dJ}{dt} = \frac{1}{16\pi} \sum_{\ell,m} i m \frac{(\ell+2)!}{(\ell-2)!} \left[\frac{d\Psi_{\ell m}^{(e)}}{dt} (\Psi_{\ell m}^{(e)})^* + \frac{d\Psi_{\ell m}^{(o)}}{dt} (\Psi_{\ell m}^{(o)})^* \right], \quad (93)$$

or, using the Moncrief master function (26) for the odd-parity perturbations [62]

$$\frac{dJ}{dt} = \frac{1}{16\pi} \sum_{\ell,m} i m \frac{(\ell+2)!}{(\ell-2)!} \left[\frac{d\Psi_{\ell m}^{(e)}}{dt} (\Psi_{\ell m}^{(e)})^* + Q_{\ell m}^{(o)} \int_{-\infty}^t (Q_{\ell m}^{(o)})^* (t') dt' \right]. \quad (94)$$

To conclude, we report the expression for the energy spectrum $dE/d\omega$, which is readily calculated from equation (88) after performing the Fourier transform of the odd and even-parity master functions, i.e.

$$\frac{dE}{d\omega} = \frac{1}{16\pi^2} \sum_{\ell,m} \frac{(\ell+2)!}{(\ell-2)!} \omega^2 \left(|\tilde{\Psi}_{\ell m}^{(e)}|^2 + |\tilde{\Psi}_{\ell m}^{(o)}|^2 \right), \quad (95)$$

where we have indicated with $\tilde{f}(\omega, r)$ the Fourier transform of the timeseries $f(t, r)$. As for the emitted power, also the energy spectrum will have a different expression if the odd-parity Moncrief function is used and in this case one obtains

$$\frac{dE}{d\omega} = \frac{1}{16\pi^2} \sum_{\ell, m} \frac{(\ell+2)!}{(\ell-2)!} \left(\omega^2 |\tilde{\Psi}_{\ell m}^{(e)}|^2 + |\tilde{Q}_{\ell m}^{(o)}|^2 \right). \quad (96)$$

We note that expression (96) for the energy spectrum agrees with the ones given in [53, 56] for the cases of odd and even-parity perturbations, respectively and with the complete expression in [52] for both odd and even-parity multipoles. On the other hand, the energy spectrum reported in [63] suffers of the same inconsistencies discussed above for equation (84), so that equation (27) in [63] should effectively be replaced by equation (96).

We conclude this section by noting that a new formalism has been recently proposed to compute the energy and angular momentum fluxes absorbed by a Schwarzschild (and Kerr) black hole in a form which is suitable for time-domain computations [62, 49]. Although the effect of the absorbed flux is generally small in astrophysical situations, it can be important in the secular dynamics in black-hole spacetimes, as discussed in [52]. Referring the interested reader to [62, 49] for more details, here we simply recall that the calculation of the absorbed energy flux cannot be made with the present choice of Schwarzschild coordinates (in which the ingoing flux at the horizon is suppressed by the infinite redshift) and must therefore be performed using an advanced time-coordinate. Furthermore, Isaacson's prescription for the stress-energy pseudo-tensor (85) also ceases to be valid in the vicinity of the horizon, where the short-wave approximation breaks down and an alternative approach needs to be implemented [62, 49].

6.5. A commonly used convention

We conclude this section on asymptotic expressions by discussing a rather popular choice for the gauge-invariant master functions and that has found successful application in the extraction of the gravitational-wave content of numerically simulated spacetimes [48, 34, 35, 38]. Furthermore, the convention discussed below has been implemented in the *Cactus* computational toolkit [36, 37], a diffused and freely available infrastructure for the numerical solution of the Einstein equations [71]. Numerous tests and applications of this implementation have been performed over the years and we refer the reader to [36, 37, 72, 73] for examples both in vacuum and non-vacuum spacetimes.

The reference work for this convention is the one by Abrahams and Price (1996) [48], although a similar approach for the even-parity part of the perturbations was also adopted in previous works [44, 47]. We first note that the coefficients c_0 , c_1 and c_2 introduced in [48] are related simply to the multipolar coefficients of the odd-parity part introduced in section 3. More specifically, $c_2 = -2h = h_2$, $c_0 = h_0^{(o)}$ and $c_1 = h_1^{(o)}$ and it is then easy to realise that the odd and even-parity master functions $Q_{\ell m}^\times$ and $Q_{\ell m}^+$ defined in [48] are related to the master functions discussed so far through the simple algebraic expressions

$$Q_{\ell m}^\times = \sqrt{\frac{2(\ell+2)!}{(\ell-2)!}} Q_{\ell m}^{(o)}, \quad (97)$$

$$Q_{\ell m}^+ = \sqrt{\frac{2(\ell+2)!}{(\ell-2)!}} \Psi_{\ell m}^{(e)}, \quad (98)$$

so that the asymptotic expression for the gravitational-wave amplitudes in the two

polarizations are given by

$$h_+ = \frac{1}{\sqrt{2}r} \sum_{\ell,m} \sqrt{\frac{(\ell-2)!}{(\ell+2)!}} \left[Q_{\ell m}^+ W^{\ell m} - \left(\int_{-\infty}^t Q_{\ell m}^\times(t') dt' \right) \frac{X^{\ell m}}{\sin \theta} \right] + \mathcal{O}\left(\frac{1}{r^2}\right), \quad (99)$$

$$h_\times = \frac{1}{\sqrt{2}r} \sum_{\ell,m} \sqrt{\frac{(\ell-2)!}{(\ell+2)!}} \left[Q_{\ell m}^+ \frac{X^{\ell m}}{\sin \theta} + \left(\int_{-\infty}^t Q_{\ell m}^\times(t') dt' \right) W^{\ell m} \right] + \mathcal{O}\left(\frac{1}{r^2}\right). \quad (100)$$

Also expressions (99) and (100) can be combined into a single one

$$h_+ - i h_\times = \frac{1}{\sqrt{2}r} \sum_{\ell,m} \left(Q_{\ell m}^+ - i \int_{-\infty}^t Q_{\ell m}^\times(t') dt' \right) {}_{-2}Y^{\ell m}(\theta, \phi) + \mathcal{O}\left(\frac{1}{r^2}\right), \quad (101)$$

which closely resembles expression (84) and that in its compactness highlights the advantage of the normalisation (97)–(98). Also very compact is the expression for the emitted power that, with this convention, simply reads

$$\frac{dE}{dt} = \frac{1}{32\pi} \sum_{\ell,m} \left(\left| \frac{dQ_{\ell m}^+}{dt} \right|^2 + |Q_{\ell m}^\times|^2 \right). \quad (102)$$

We note that expression (102) corrects equation (24) of [48], where a time derivative of the odd-parity perturbations is present, leading to an obvious dimensional inconsistency. This typo was subsequently corrected in [53].

7. Conclusions

Black hole perturbation theory has been and will continue to be a powerful tool with continuous developments and multiple applications. In this paper we have briefly reviewed the theory of metric perturbations of Schwarzschild black hole, concentrating in particular on its gauge-invariant formulations. Special care has been paid to “filter” the technical details in the mathematical apparatus that may obscure the important results and to “steer” the reader through the numerous conventions and notations that have accompanied the development of the formalism over the years, pointing out the misprints and errors that can be found in the literature. Among the aims of this review is that of providing the reader with a set of expressions in the two most common formulations and that can be readily used for the calculation of the odd and even-parity perturbations of a Schwarzschild spacetime in the presence of generic matter-sources. In addition, we have reported the asymptotic expressions for the gravitational-wave amplitudes and losses driven by these generic perturbations. The latter can find useful application in the calculation of the gravitational-wave content of numerically simulated spacetimes.

With suitable gauge choices, much of what presented here for a Schwarzschild black hole could be extended to a Kerr background and this will be the subject of future work.

Acknowledgments

It is a pleasure to thank Andrew Abrahams, Tomohiro Harada, Frank Herrmann, Vincent Moncrief, Eric Poisson, Richard Price, Ed Seidel and Manuel Tiglio for useful discussions and comments. The activity of AN at the University of Valencia was supported by a “Angelo della Riccia” fellowship.

Appendix A. Some explicit expressions

In the general expressions given in the main text the tensor harmonic appear in terms of covariant derivatives. While compact, such expressions need to be ‘unwrapped’ in practice and in what follows we provide their explicit expressions.

$$\nabla_\theta \nabla_\theta Y^{\ell m} = \partial_\theta^2 Y^{\ell m} , \quad (\text{A.1})$$

$$\nabla_\phi \nabla_\phi Y^{\ell m} = \partial_\phi^2 Y^{\ell m} + \sin \theta \cos \theta \partial_\theta Y^{\ell m} , \quad (\text{A.2})$$

$$\nabla_\phi \nabla_\theta Y^{\ell m} = \partial_{\theta\phi}^2 Y^{\ell m} - \cot \theta \partial_\phi Y^{\ell m} , \quad (\text{A.3})$$

for the even-parity ones, and

$$S_\theta^{\ell m} = -\frac{1}{\sin \theta} \partial_\phi Y^{\ell m} , \quad S_\phi^{\ell m} = \sin \theta \partial_\theta Y^{\ell m} , \quad (\text{A.4})$$

$$\nabla_\theta S_\theta^{\ell m} = -\frac{1}{\sin \theta} (\partial_\theta^2 Y^{\ell m} - \cot \theta \partial_\phi Y^{\ell m}) , \quad (\text{A.5})$$

$$\nabla_\phi S_\phi^{\ell m} = \sin \theta (\partial_\phi^2 Y^{\ell m} - \cot \theta \partial_\theta Y^{\ell m}) , \quad (\text{A.6})$$

$$\nabla_\theta S_\phi^{\ell m} = \sin \theta \partial_\theta^2 Y^{\ell m} , \quad (\text{A.7})$$

$$\nabla_\phi S_\theta^{\ell m} = -\frac{1}{\sin \theta} \partial_\phi^2 Y^{\ell m} - \cos \theta \partial_\theta Y^{\ell m} . \quad (\text{A.8})$$

for the vector and tensor odd-parity part.

We also recall that the orthogonality relations between scalar, vector and tensor harmonics imply

$$\int d\Omega Y_{\ell m}^* Y_{\ell' m'} = \delta_{\ell\ell'} \delta_{mm'} , \quad (\text{A.9})$$

$$\int d\Omega \gamma^{cd} (\partial_c Y_{\ell m}^*) (\partial_d Y_{\ell' m'}) = \Lambda \delta_{\ell\ell'} \delta_{mm'} , \quad (\text{A.10})$$

$$\int d\Omega Z_{cd}^{*\ell m} Z_{\ell m}^{cd} = \frac{\Lambda(\Lambda-2)}{2} \delta_{\ell\ell'} \delta_{mm'} , \quad (\text{A.11})$$

while the orthogonality relations between the odd-parity axial vectors S_c and tensor $\nabla_{(c} S_{d)}$ follow straightforwardly from the above equations, giving

$$\int d\Omega \gamma^{cd} S_c^{*\ell m} S_d^{\ell' m'} = \Lambda \delta_{\ell\ell'} \delta_{mm'} , \quad (\text{A.12})$$

$$\int d\Omega \nabla_{(c} S_{d)}^{*\ell m} \nabla^{(c} S_{\ell' m')}^d = 2\Lambda(\Lambda-1) \delta_{\ell\ell'} \delta_{mm'} . \quad (\text{A.13})$$

Appendix B. A useful example: point-particles

In the spirit of providing a useful example of a matter-source generating odd and even-parity perturbations of a Schwarzschild spacetime, we now a particular problem that has been carefully studied over the years: that of a point-particle orbiting around a black hole. Since this problem has been solved in many different cases and using very different approaches, it can serve as a useful test for codes handling more complex matter-sources. Because of this, we report below the relevant components of the odd and even-parity decomposition of the particle’s energy-momentum tensor.

For a particle of mass $\mu \ll M$, specific energy E and angular momentum L , the equations of motion in the equatorial plane ($\theta = \pi/2$) for the coordinates $R(t)$ and $\Phi(t)$

are

$$\dot{R}^2 = \left(1 - \frac{2M}{R}\right)^2 \left[1 - \frac{1}{E^2} \left(1 - \frac{2M}{R}\right) \left(1 + \frac{L^2}{R^2}\right)\right], \quad (\text{B.1})$$

$$\dot{\Phi} = \left(1 - \frac{2M}{R}\right) \frac{L}{R^2 E}, \quad (\text{B.2})$$

where $\dot{R} \equiv dR/dt$, $\dot{\Phi} \equiv d\Phi/dt$ and where the energy-momentum tensor reads

$$t_{\mu\nu} = \frac{\mu}{r^2 E} \left(1 - \frac{2M}{r}\right) \delta(r - R(t)) \delta(\phi - \Phi(t)) \delta\left(\theta - \frac{\pi}{2}\right) \times \quad (\text{B.3})$$

$$\times \begin{pmatrix} E^2 & E^2 (1 - 2M/r)^{-2} \dot{R} & 0 & -LE \\ \text{sym} & E^2 (1 - 2M/r)^{-4} \dot{R}^2 & 0 & LE (1 - 2M/r)^{-2} \dot{R} \\ \text{sym} & \text{sym} & 0 & 0 \\ \text{sym} & \text{sym} & \text{sym} & L^2 \end{pmatrix}. \quad (\text{B.4})$$

As a result, the odd-parity components of the stress energy tensor are[¶]

$$L^{\ell m} = -\frac{im}{\Lambda(\Lambda - 1)} \frac{\mu L^2}{r^2 E} \left(1 - \frac{2M}{r}\right) \delta(r - R(t)) \partial_\theta Y_{\ell m}^*, \quad (\text{B.5})$$

$$L_0^{\ell m} = -\frac{1}{\Lambda} \frac{\mu L}{r^2} \left(1 - \frac{2M}{r}\right) \delta(r - R(t)) \partial_\theta Y_{\ell m}^*, \quad (\text{B.6})$$

$$L_1^{\ell m} = \frac{1}{\Lambda} \frac{\mu L}{r^2} \left(1 - \frac{2M}{r}\right)^{-1} \dot{R} \delta(r - R(t)) \partial_\theta Y_{\ell m}^*, \quad (\text{B.7})$$

while the even-parity components assume the form

$$T_{00}^{\ell m} = \mu E \frac{r - 2M}{r^3} \delta(r - R(t)) Y_{\ell m}^*, \quad (\text{B.8})$$

$$T_{11}^{\ell m} = \frac{\mu E r}{(r - 2M)^3} (\dot{R})^2 \delta(r - R(t)) Y_{\ell m}^*, \quad (\text{B.9})$$

$$T_{01}^{\ell m} = -\frac{\mu}{r(r - 2M)} \dot{R} \delta(r - R(t)) Y_{\ell m}^*, \quad (\text{B.10})$$

$$T_0^{\ell m} = \frac{im\mu L}{\Lambda} \frac{r - 2M}{r^3} \delta(r - R(t)) Y_{\ell m}^*, \quad (\text{B.11})$$

$$T_1^{\ell m} = -im \frac{\mu L}{\Lambda r(r - 2M)} \dot{R} \delta(r - R(t)) Y_{\ell m}^*, \quad (\text{B.12})$$

$$T_2^{\ell m} = \frac{\Lambda - 2m^2}{\Lambda(\Lambda - 2)} \frac{\mu L^2 (r - 2M)}{E r^3} \delta(r - R(t)) Y_{\ell m}^*, \quad (\text{B.13})$$

$$T_3^{\ell m} = \frac{\mu L^2 (r - 2M)}{2E r^5} \delta(r - R(t)) Y_{\ell m}^*. \quad (\text{B.14})$$

Appendix C. Variational principle for odd-parity perturbations

In Sect. 4 we have discussed that it is possible to choose two different representations for the odd-parity perturbations, namely expressions (20) for the CPM convention and expression

[¶] Note that the spherical harmonics and their derivatives are all evaluated at $\phi = \Phi(t)$ and $\theta = \pi/2$.

(26) for the RW convention. An intimate connection links the two master functions and this can be appreciated by using the variational formalism employed by Moncrief in [14]. To highlight this, let us recall that two perturbation functions k_1, k_2 can be introduced for the odd-parity perturbations

$$k_1 \equiv h_1^{(o)} + \frac{1}{2} \left(\partial_r h_2 - \frac{2}{r} h_2 \right), \quad k_2 \equiv h_2 = -2h, \quad (\text{C.1})$$

where k_1 is gauge-invariant (cf equation (18)) but k_2 is not. While this may seem just another possible choice for the odd-parity perturbations, it points out that it is now possible to introduce the quantities π_1 and π_2 as conjugate momenta of k_1 and k_2 in the Hamiltonian

$$\begin{aligned} H_T \equiv \int dr \mathcal{H} = & \frac{1}{\Lambda} \int dr \left[\frac{1}{2} \pi_1^2 + \frac{2r(r-2M)}{(\Lambda-2)} \left(\pi_2 - \frac{1}{2} \partial_r \pi_1 - \frac{1}{r} \pi_1 \right)^2 \right] \\ & + \frac{1}{2} \Lambda \int dr \left[\frac{\Lambda-2}{r^2} \left(1 - \frac{2M}{r} \right) k_1^2 \right] - 2 \int dr h_0^{(o)} \pi_2. \end{aligned} \quad (\text{C.2})$$

In addition to the constraint equation $\pi_2 = 0$ (which must be enforced at initial time and is then conserved since k_2 is cyclic), the other Hamilton equations are

$$\partial_t k_1 = \frac{\delta H_T}{\delta \pi_1}, \quad \partial_t k_2 = \frac{\delta H_T}{\delta \pi_2}, \quad \partial_t \pi_1 = -\frac{\delta H_T}{\delta k_1}, \quad (\text{C.3})$$

where $\delta(\)/\delta(\)$ refers to a functional differentiation. The first and third of eqs. (C.3) have explicit expressions

$$\partial_t k_1 = \frac{1}{\Lambda(\Lambda-2)} \left[(\Lambda-2)\pi_1 - \left(1 - \frac{2M}{r} \right) \partial_r^2 (r^2 \pi_1) + \frac{2}{r} \left(1 - \frac{3M}{r} \right) \partial_r (r^2 \pi_1) \right], \quad (\text{C.4})$$

$$\partial_t \pi_1 = -\frac{\Lambda(\Lambda-2)}{r^2} \left(1 - \frac{2M}{r} \right) k_1. \quad (\text{C.5})$$

and the Regge-Wheeler equation in the RWM convention can be obtained by time-differentiating equation (C.4), replacing $\partial_t \pi_1$ through equation (C.5) and introducing the quantity

$$Q^{(o)} = \frac{k_1}{r} \left(1 - \frac{2M}{r} \right), \quad (\text{C.6})$$

to obtain

$$\partial_t^2 Q^{(o)} - \partial_{r_*}^2 Q^{(o)} + \left(1 - \frac{2M}{r} \right) \left(\frac{\Lambda}{r^2} - \frac{6M}{r^3} \right) Q^{(o)} = 0. \quad (\text{C.7})$$

Equation (C.7) coincides with equation (4.20) of [14] with $Q = Q^{(o)}$.

Remarkably, the Regge-Wheeler equation in the CPM convention can be obtained also through the Hamilton equations in terms of the conjugate moment to k_1 , i.e. $r\pi_1$. More specifically, it is sufficient to differentiate in time equation (C.5) and to replace $\partial_t k_1$ in equation (C.4) to obtain

$$\partial_t^2 (r\pi_1) - \partial_{r_*}^2 (r\pi_1) + \left(1 - \frac{2M}{r} \right) \left(\frac{\Lambda}{r^2} - \frac{6M}{r^3} \right) r\pi_1 = 0, \quad (\text{C.8})$$

Equation (C.7) coincides with equation (II-15) of [15] with

$$r\pi_1 = \frac{\tilde{\psi}}{\Lambda} \equiv r \left[\partial_t h_1^{(o)} - r^2 \partial_r \left(\frac{h_0^{(o)}}{r^2} \right) \right]. \quad (\text{C.9})$$

Note also that the second of Hamilton equations (C.3) can be used to derive

$$\partial_t h_2 = -\frac{2}{\Lambda(\Lambda-2)} \left(1 - \frac{2M}{r}\right) \partial_r (r^2 \pi_1) - 2h_0^{(o)}, \quad (\text{C.10})$$

which coincides with equation (III-20) of Cunningham *et al.* [15] and has been used to obtain the asymptotic relation between $\Psi^{(o)}$ and the gravitational wave amplitude (cf equation (60)).

Finally, we underline that the relation (27) between the two master functions $\Psi^{(o)}$ and $Q^{(o)}$ can be obtained in coordinate independent form also in the formalism introduced by Martín-García and Gundlach [39]. More specifically, exploiting the odd-parity perturbations equations (47) as

$${}^0 g^{CD} \nabla_D [4r(\nabla_{[A} r)k_{C]} + 2r^2 \nabla_{[C} k_{A]}] n^A - (\Lambda - 2)k_A n^A = -16\pi L_A^{\ell m} n^A \quad (\text{C.11})$$

and the gauge-invariant and coordinate independent definitions of $\Psi^{(o)}$ and $Q^{(o)}$ (i.e. equations (19) and (26) of the main text), straightforward calculations then yield

$$\dot{\Psi}^{(o)} = -\frac{Q^{(o)}}{r'} + \frac{16\pi}{\Lambda-2} r n^A L_A^{\ell m} \quad (\text{C.12})$$

where $\dot{\Psi}^{(o)} \equiv u^A \nabla_A \Psi^{(o)}$ and $r' \equiv n^A \nabla_A r$ are ‘frame derivatives’ in terms of the non-coordinate basis vectors (u^A, n^A) of M^2 , such that $-u_A u^A = n_A n^A = 1$, ${}^0 g_{AB} = -u_A u_B + n_A n_B$ and $\epsilon_{AB} = n_A u_B - u_A n_B$. In the case of Schwarzschild coordinates, $n^A = (0, e^{-b})$, $u^A = (e^b, 0)$ and equation (C.12) coincides with equation (27).

Appendix D. Angular pattern functions

Finally, this section is devoted to a list of the expressions of the angular functions $W^{\ell m}$ and $X^{\ell m}$ for the first values of ℓ and m . While these expressions can be easily reproduced with straightforward algebra, they are tedious to derive and hard to find in the literature.

- $\ell = 2$

$$W^{20} = \sqrt{\frac{45}{4\pi}} \sin^2 \theta, \quad W^{21} = e^{i\phi} \sqrt{\frac{15}{8\pi}} \sin 2\theta, \quad W^{22} = e^{i2\phi} \sqrt{\frac{15}{32\pi}} (3 + \cos 2\theta) \quad (\text{D.1})$$

$$X^{20} = 0, \quad X^{21} = ie^{i\phi} \sqrt{\frac{15}{2\pi}} \sin^2 \theta, \quad X^{22} = ie^{i2\phi} \sqrt{\frac{15}{8\pi}} \sin 2\theta. \quad (\text{D.2})$$

- $\ell = 3$

$$W^{30} = \frac{15}{2} \sqrt{\frac{7}{\pi}} \cos \theta \sin^2 \theta, \quad W^{31} = e^{i\phi} \frac{5}{8} \sqrt{\frac{21}{\pi}} (1 + 3 \cos 2\theta) \sin \theta, \quad (\text{D.3})$$

$$W^{32} = e^{i2\phi} \sqrt{\frac{105}{32\pi}} (1 + 3 \cos 2\theta) \cos \theta, \quad W^{33} = -e^{i3\phi} \sqrt{\frac{315}{64\pi}} (3 + \cos 2\theta) \sin \theta, \quad (\text{D.4})$$

$$X^{31} = ie^{i\phi} \sqrt{\frac{525}{4\pi}} \cos \theta \sin^2 \theta, \quad X^{32} = ie^{i2\phi} \sqrt{\frac{105}{2\pi}} \sin \theta \cos 2\theta, \quad (\text{D.5})$$

$$X^{33} = -ie^{i3\phi} \frac{3}{2} \sqrt{\frac{35}{\pi}} \cos \theta \sin^2 \theta. \quad (\text{D.6})$$

- $\ell = 4$

$$W^{40} = \frac{45}{8\sqrt{\pi}} (5 + 7 \cos 2\theta) \sin^2 \theta, \quad W^{41} = e^{i\phi} \frac{9}{8} \sqrt{\frac{5}{\pi}} (7 \cos 2\theta - 1) \sin 2\theta, \quad (\text{D.7})$$

$$W^{42} = e^{i2\phi} \frac{9}{16} \sqrt{\frac{5}{2\pi}} (5 + 4 \cos 2\theta + 7 \cos 4\theta), \quad W^{43} = -ie^{i3\phi} \frac{9}{2} \sqrt{\frac{35}{\pi}} \sin \theta \cos^3 \theta, \quad (\text{D.8})$$

$$W^{44} = e^{i4\phi} \frac{9}{8} \sqrt{\frac{35}{2\pi}} (3 + \cos 2\theta) \sin^2 \theta, \quad X^{41} = ie^{i\phi} \frac{9}{8} \sqrt{\frac{5}{\pi}} (7 \cos 2\theta + 5) \sin^2 \theta, \quad (\text{D.9})$$

$$X^{42} = ie^{i2\phi} \frac{9}{8} \sqrt{\frac{5}{2\pi}} (7 \cos 2\theta - 3) \sin 2\theta, \quad X^{43} = -ie^{i3\phi} \frac{9}{8} \sqrt{\frac{35}{\pi}} (1 + 3 \cos 2\theta) \sin^2 \theta, \quad (\text{D.10})$$

$$X^{44} = ie^{i4\phi} \frac{9}{2} \sqrt{\frac{35}{2\pi}} \cos \theta \sin^2 \theta. \quad (\text{D.11})$$

- $\ell = 5$

$$W^{50} = \frac{105}{16} \sqrt{\frac{11}{\pi}} (5 \cos \theta + 3 \cos 3\theta) \sin^2 \theta, \quad (\text{D.12})$$

$$W^{51} = e^{i\phi} \frac{7}{32} \sqrt{\frac{165}{2\pi}} (5 + 12 \cos 2\theta + 15 \cos 4\theta) \sin \theta, \quad (\text{D.13})$$

$$W^{52} = e^{i2\phi} \sqrt{\frac{1155}{512\pi}} (13 - 12 \cos 2\theta + 15 \cos 4\theta) \cos \theta, \quad (\text{D.14})$$

$$W^{53} = -e^{i3\phi} \frac{3}{64} \sqrt{\frac{385}{\pi}} (21 + 28 \cos 2\theta + 15 \cos 4\theta) \sin \theta, \quad (\text{D.15})$$

$$W^{54} = e^{i4\phi} 3 \sqrt{\frac{385}{128\pi}} (7 + 5 \cos 2\theta) \cos \theta \sin^2 \theta, \quad (\text{D.16})$$

$$W^{55} = -e^{i5\phi} \frac{15}{16} \sqrt{\frac{77}{\pi}} (3 + \cos 2\theta) \sin^3 \theta, \quad (\text{D.17})$$

$$X^{50} = 0, \quad (\text{D.18})$$

$$X^{51} = ie^{i\phi} \frac{7}{4} \sqrt{\frac{165}{2\pi}} (1 + 3 \cos 2\theta) \sin^2 \theta, \quad (\text{D.19})$$

$$X^{52} = ie^{i2\phi} \sqrt{\frac{1155}{32\pi}} (\cos 2\theta + 3 \cos 4\theta) \sin \theta, \quad (\text{D.20})$$

$$X^{53} = -ie^{i3\phi} 3 \sqrt{\frac{385}{64\pi}} (9 \cos 2\theta - 1) \cos \theta \sin^2 \theta, \quad (\text{D.21})$$

$$X^{54} = ie^{i4\phi} 3 \sqrt{\frac{385}{8\pi}} (1 + 2 \cos 2\theta) \sin^3 \theta, \quad (\text{D.22})$$

$$X^{55} = -ie^{i5\phi} 15 \sqrt{\frac{77}{16\pi}} \cos \theta \sin^4 \theta. \quad (\text{D.23})$$

References

- [1] S. Chandrasekhar, *The Mathematical Theory of Black Holes*, Oxford University Press, New York (1983).
- [2] V.P. Frolov and I.D. Novikov, *Black Hole Physics*, Kluwer Academic Publishers, Dordrecht (1998).

- [3] K.D. Kokkotas and B.G. Schmidt, Living Rev. Relativ. **2**, 2 (1999).
- [4] T. Regge and J. A. Wheeler, Phys. Rev. **108**, 1063 (1957).
- [5] R. H. Price, Phys. Rev. D **5**, 2419 (1972).
- [6] R. H. Price, Phys. Rev. D **5**, 2439 (1972).
- [7] C. V. Vishveshwara, Phys. Rev. D **1**, 2870 (1970).
- [8] C. V. Vishveshwara, Nature **227**, 936 (1970).
- [9] W. H. Press, Astrophys. J. Lett. **170**, L105 (1971).
- [10] S. Chandrasekhar and S. Detweiler, Proc. R. Soc. London **A344**, 441 (1975).
- [11] C. W. Misner, K. S. Thorne and J. A. Wheeler, *Gravitation*, Freeman, New York (1973).
- [12] F. J. Zerilli, Phys. Rev. Lett. **24**, 737 (1970).
- [13] F. J. Zerilli, Phys. Rev. D **2**, 2141 (1970).
- [14] V. Moncrief, Annals of Physics, **88**, 323 (1974).
- [15] C. T. Cunningham, R. H. Price, and V. Moncrief, Astrophys. J. **224**, 643 (1978).
- [16] C. T. Cunningham, R. H. Price, and V. Moncrief, Astrophys. J. **230**, 870 (1979).
- [17] S. A. Teukolsky, Phys. Rev. Lett. **29**, 1114 (1972).
- [18] S. A. Teukolsky, Astrophys. J. **185**, 635 (1973).
- [19] U. H. Gerlach and U. K. Sengupta, Phys. Rev. D **19**, 2268 (1979).
- [20] U. H. Gerlach and U. K. Sengupta, J. Math. Phys. **20**, 2540 (1979).
- [21] U. H. Gerlach and U. K. Sengupta, Phys. Rev. D **22**, 1300 (1980).
- [22] M. Karlovini, Class. Q. Grav. **19**, 2125 (2002).
- [23] E. Seidel and T. Moore, Phys. Rev. D **35**, 2287 (1987).
- [24] E. Seidel, E. S. Myra and T. Moore, Phys. Rev. D **38**, 2349 (1988).
- [25] E. Seidel, Phys. Rev. D **42**, 1884 (1990).
- [26] E. Seidel, Phys. Rev. D **44**, 950 (1991).
- [27] H. Kodama, A. Ishibashi and O. Seto, Phys. Rev. D **62**, 064022 (2000).
- [28] H. Kodama and A. Ishibashi, Prog. Theor. Phys. **110**, 701 (2003).
- [29] A. Ishibashi and H. Kodama, Prog. Theor. Phys. **110**, 901 (2003).
- [30] H. Kodama and A. Ishibashi, Prog. Theor. Phys. **111**, 29 (2004).
- [31] R. H. Price and J. Pullin, Phys. Rev. Lett. **72**, 3297 (1994).
- [32] A. M. Abrahams and G. Cook, Phys. Rev. D **50**, 2364 (1994).
- [33] A. M. Abrahams, S. L. Shapiro and S. A. Teukolsky, Phys. Rev. D **51**, 4295 (1995).
- [34] A. M. Abrahams, L. Rezzolla, M. E. Rupright, et al. Phys. Rev. Lett. **80** 1812 (1998).
- [35] M.E. Rupright, A.M. Abrahams, L. Rezzolla, Phys. Rev. D **58**, 044005 (1998).
- [36] K. Camarda, E. Seidel, Phys. Rev. D **59**, 064019 (1999).
- [37] G. Allen, K. Camarda, E. Seidel, gr-qc/9806036.
- [38] L. Rezzolla, A. M. Abrahams, R. A. Matzner, M. Rupright, S. L. Shapiro, Phys. Rev. D **59**, 064001 (1999).
- [39] J. M. Martín-García and C. Gundlach, Phys. Rev. D **59**, 064031 (1999).
- [40] C. Gundlach and J. M. Martín-García Phys. Rev. D **61**, 084024 (2000).
- [41] C. Gundlach and J. M. Martín-García, Phys. Rev. D **64**, 024012 (2001).
- [42] F. J. Zerilli, J. Math. Phys. **11**, 2203 (1970).
- [43] J. Mathews, J. Soc. Ind. Appl. Math. **10**, 768 (1962).
- [44] A. Abrahams, D. Bernstein, D. Hobill, E. Seidel and L. Smarr, Phys. Rev. D **45**, 3544 (1992).
- [45] P. Anninos, D. Hobill, E. Seidel, L. Smarr and W.M. Suen, Phys. Rev. Lett. **71**, 2851 (1993).
- [46] P. Anninos, D. Hobill, E. Seidel, L. Smarr and W.M. Suen, Phys. Rev. D **52**, 2044 (1995).
- [47] P. Anninos, R. H. Price, J. Pullin, E. Seidel and W.M. Suen, Phys. Rev. D **52**, 4462 (1995).
- [48] A. Abrahams, R. H. Price, Phys. Rev. D **53**, 1963 (1996).; **53** 1972 (1996).
- [49] K. Martel and E. Poisson, Phys. Rev. D **71**, 104003 (2005).
- [50] T. Harada, H. Iguchi and M. Shibata, Phys. Rev. D **68**, 024002 (2003).
- [51] O. Sarbach and M. Tiglio, Phys. Rev. D **64**, 084016 (2001).
- [52] K. Martel, Phys. Rev. D **69**, 044025 (2004).
- [53] Z. Andrade and R. H. Price, Phys. Rev. D **60**, 104037 (1999).
- [54] K. Tominaga, M. Saijo, and K. Maeda, Phys. Rev. D **60**, 024004 (1999).
- [55] V. Ferrari and K. D. Kokkotas, Phys. Rev. D **62**, 107504 (2000).
- [56] C. O. Lousto and R. H. Price, Phys. Rev. D **55**, 2124 (1997).
- [57] J. Ruoff, Phys. Rev. D **63**, 064018 (2001).
- [58] J. Ruoff, P. Laguna and J. Pullin, Phys. Rev. D **63**, 064019 (2001).
- [59] K. Martel and E. Poisson, Phys. Rev. D **66**, 084001 (2002).
- [60] A. Nagar, G. Diaz, J.A. Pons, and J.A. Font, Phys. Rev. D **69**, 124028 (2004).
- [61] A. Nagar, J.A. Font, O. Zanotti, and R. De Pietri, Phys. Rev. D **71** in press (2005).
- [62] E. Poisson, Phys. Rev. D **70**, 084044 (2004).
- [63] M. Kawamura and K. Oohara, Prog. Theor. Phys. **111**, 589 (2004).

- [64] M. Kawamura, K. Oohara and T. Nakamura, [astro-ph/0306481](#).
- [65] J.M. Bardeen and W.H. Press, *J. Math. Phys.* **14**, 7 (1973).
- [66] M. Shibata, K. Taniguchi, and K. Uryū, *Phys. Rev. D* **68**, 084020 (2003).
- [67] M. Shibata and Y. Sekiguchi, *Phys. Rev. D* **68**, 104020 (2003).
- [68] M. Shibata and Y. Sekiguchi, *Phys. Rev. D* **71** *in press* 024014 (2005).
- [69] R. Isaacson, *Phys. Rev.* **166**, 1263 (1968).; *ibid.* **166**, 1272 (1968).
- [70] L. D. Landau, and E. M. Lifshitz: *The Classical Theory Of Fields*, 4th ed., London: Pergamon Press (1975).
- [71] Cactus Computational Toolkit, www.cactuscode.org.
- [72] J. A. Font, *et al.* *Phys. Rev. D* 65, 084024 (2002).
- [73] L. Baiotti, I. Hawke, L. Rezzolla, and E. Schnetter, *Phys. Rev. Lett.* **94**, 131101 (2005).